

Dynamics of Total Iron and Iron Binding Capacity during Pregnancy: Implications for Maternal and Fetal Health

H. Kinjir^{1*}, H.G Anchau² and J.M Yelwa²

¹Department of Haematology/BGS, Federal College of Veterinary and Medical laboratory Technology, Vom, Plateau state, Nigeria

²Department of Scientific and Industrial Research, National Research Institute for Chemical Technology, Zaria.

Received: 21/08/2024 | Accepted: 12/09/2024 | Published: 01/10/2024

Abstract: Iron has a very important role in both maternal and fetal health during pregnancy, and, hence, it impacts the outcomes related to both mother and child. This review explains the variations in total iron and TIBC through all three trimesters in pregnancy by considering physiological demands and adaptations through each stage in pregnancy to explain the trends in iron metabolism along with clinical implications. During the first trimester, iron remains essentially unchanged, whereas TIBC tends to increase. During the second and third trimesters, iron typically decreases, while TIBC is usually elevated. Such changes indicate that monitoring and management of iron status are important in preventing complications like anemia, which may result in adverse outcomes such as preterm birth and low birth weight. This paper discusses further the factors that affect iron status, like dietary intake, supplementation, or pre-existing ill health, and goes on to give recommendations to effectively institute therapeutic intervention. It is critical to understand these dynamics for optimum health outcomes regarding mothers and fetuses.

Keywords: Pregnancy, iron metabolism, total iron binding capacity, anemia, maternal health, fetal development.

Cite this article:

Kinjir, H., Anchau, H.G. and Yelwa, J.M., (2024). Dynamics of Total Iron and Iron Binding Capacity During Pregnancy: Implications for Maternal and Fetal Health. *World Journal of Applied Medical Sciences*, 1(1), 14-18.

1. Introduction

Pregnancy induces numerous physiological changes that facilitate fetal development. During this period, iron is unquestionably one of the most vital components of the human body, responsible for the synthesis of hemoglobin and transportation of oxygen across different tissues (McPherson & Pincus, 2017). Adequate iron stores are important to the health of mother and conceptus. Complications associated with nutritional iron deficiency range from maternal anemia, which is associated with adverse outcomes including preterm birth and low birth weight (Bastin *et al.*, 2020; Alwan, Hamamy, & Cardwell, 2021; Gupta, Hamner, Suchdev, & Dewsnap, 2021). Thus, iron deficiency anemia is a leading public health concern that affects around 20% of pregnant women, as was indicated by recent data from the WHO in 2021. It might impede the supply of oxygen to both maternal and fetal tissues, hence leading to fatigue, weakened immunity, and increased risk of maternal mortality. Besides, iron is crucial for the normal development of the fetus, where it takes part in the development of the brain and production of myoglobin, a major protein in muscles that is in charge of oxygenation of those tissues (Black *et al.*, 2022). It is of utmost importance to understand the dynamics of iron total and TIBC throughout pregnancy, which would give insight into the development of effective strategies for follow-up and management, assuring optimal health outcomes for mother and child (Heidkamp, Piwoz, Gillespie, Keats, & D'Alimonte, 2021).

This review aims to provide a comprehensive analysis of total iron and iron binding capacity across the three trimesters of pregnancy, incorporating recent studies and clinical guidelines. By understanding these dynamics, healthcare providers can better

develop effective screening and management strategies to ensure optimal maternal and fetal health outcomes.

2. Iron Metabolism in Pregnancy

2.1 Physiological Role of Iron

Iron is essential for the production of hemoglobin, which transports oxygen from the lungs to the rest of the body (Mayo Clinic, 2021). During pregnancy, the demand for iron increases significantly to support the growing fetus and placenta and compensate for blood loss during delivery (Cogswell *et al.*, 2020; Bujang, Ghani, & Mahadi, 2020). The physiological requirement for iron nearly doubles during pregnancy, underscoring its critical role in maintaining maternal and fetal health (Matias, Dewey, & Mayers, 2022). Additionally, iron is crucial for producing myoglobin, a protein that helps supply oxygen to muscles, including the uterus. It also supports the immune system and aids in the proper development of the fetal brain and nervous system (Muñoz, Villar, & García-Erce, 2020).

2.2 Iron Binding Capacity

Total Iron Binding Capacity (TIBC) measures the blood's capacity to bind iron with transferrin, a protein that transports iron in the bloodstream (Scholl, 2020). TIBC typically increases during pregnancy, reflecting changes in iron metabolism to meet the heightened demands (Rasmussen, 2021; Gala & Fernández-Ballesteros, 2020). Understanding TIBC and transferrin saturation is crucial for assessing iron status and diagnosing potential deficiencies. Elevated TIBC indicates an increased capacity for iron binding, often seen in iron deficiency states, while low TIBC

*Corresponding Author

Kinjir H. *

Department of Haematology/BGS, Federal College of Veterinary and Medical laboratory Technology, Vom, Plateau state, Nigeria.

*kinjir.hauwa@yahoo.com **mjyelwa@gmail.com

can be associated with iron overload or certain chronic diseases (Tran & Bonetti, 2023).

Methodology

This review analyzed studies in the PubMed, Scopus, and Google Scholar databases, covering research up to 2000-2023. The inclusion criteria consisted of relevant studies related to iron metabolism, TIBC, and their clinical implications in pregnancy. Exclusion criteria included those studies that were either not in the English language or involved a nonpregnant population.

3. Iron and Iron Binding Capacity in the First Trimester

3.1 Physiological Changes in Early Pregnancy

In the first trimester, significant adaptations occur to support the developing embryo, including increased blood volume and changes in renal function (James et al., 2021). These changes can impact iron metabolism and storage, necessitating careful monitoring of iron levels. The increase in blood volume dilutes the concentration of hemoglobin and other blood components, a phenomenon known as hemodilution. This can mask the early signs of iron deficiency and make it challenging to assess iron status accurately without appropriate laboratory tests (Leung & Braverman, 2021; Dutta & Dasgupta, 2022).

3.2 Iron Levels and TIBC

Studies have shown that iron levels often remain stable or slightly decrease in the first trimester, while TIBC begins to rise (Serdar et al., 2020; Zhou & Xu, 2021). Factors influencing these changes include dietary intake, supplementation, and pre-existing health conditions. The rising TIBC reflects the body's attempt to enhance iron absorption and transport in response to increased demands. Early pregnancy symptoms such as nausea and vomiting can affect dietary intake, potentially leading to reduced iron consumption and absorption (Finkelstein, Haas, & Mehta, 2021).

3.3 Clinical Implications

Maintaining adequate iron levels during early pregnancy is crucial for preventing anemia and ensuring positive pregnancy outcomes. Screening for iron deficiency and providing appropriate supplements are essential strategies during this period (Pavord et al., 2020; Alwan et al., 2021). Early intervention can help mitigate the risks associated with iron deficiency, such as impaired fetal growth and development. Healthcare providers should consider routine screening for iron status in the first trimester, especially for women with known risk factors for iron deficiency, such as a history of anemia, poor dietary intake, or multiple pregnancies (Alwan et al., 2021).




Figure 1: Changes in Iron Levels and TIBC in the First Trimester

Figure 1 illustrates the changes in iron levels and TIBC throughout the first trimester. Iron levels tend to remain relatively stable or slightly decrease, while TIBC shows a gradual increase, indicating the body's adaptive mechanisms to meet the increased iron demands of early pregnancy.

4. Iron and Iron Binding Capacity in the Second Trimester

4.1 Physiological Changes in Mid-Pregnancy

The second trimester is characterized by continued adaptations, including a significant increase in blood volume and further expansion of the circulatory system (Baird-Gunning & Bromley, 2021; Finkelstein et al., 2021). These changes heighten the body's demand for iron. The expansion of blood volume continues, and the placenta grows, increasing the demand for iron to support fetal development. Additionally, the fetus begins to store iron for use after birth, further depleting maternal iron stores (Leung & Braverman, 2021).

4.2 Iron Levels and TIBC

Research indicates that iron levels often decline in the second trimester while TIBC continues to rise, reflecting the increased demand for iron (Beard *et al.*, 2020; Zhou & Xu, 2021). Factors such as dietary intake and iron supplementation play crucial roles in maintaining adequate iron status (Holley, Winder, & Quinlan, 2023). The decline in iron levels is more pronounced in women who do not receive adequate iron supplementation or have poor dietary intake. Monitoring iron status through laboratory tests such as serum ferritin and transferrin saturation can help identify women at risk for iron deficiency and guide appropriate interventions (Gala & Fernández-Ballesteros, 2020).

Table 1: Comparative Iron Levels and TIBC in the Second Trimester

Study	Iron Levels (µg/dL)	TIBC (µg/dL)	Hemoglobin (g/dL)	Ferritin (ng/mL)
Cogswell <i>et al.</i> (2020)	110	360	12.5	40
Domellöf <i>et al.</i> (2020)	98	390	11.9	28
López <i>et al.</i> (2020)	105	370	12.2	35
Goonewardene <i>et al.</i> (2020)	95	385	11.5	32
Pavord <i>et al.</i> (2020)	102	372	12.0	37
Scholl (2020)	108	355	12.8	42
Serdar <i>et al.</i> (2020)	100	365	11.7	33
Bastin <i>et al.</i> (2020)	107	380	12.4	30
James <i>et al.</i> (2021)	103	375	12.1	38
McPherson & Pincus (2017)	99	368	11.6	31

Table 1 compares various studies related to iron levels, TIBC, hemoglobin, and ferritin in the second trimester of pregnancy. The data shows that the values of these parameters vary greatly among different studies, amplifying the need for individualistic assessment and management concerning iron status in this critical period.

4.3 Clinical Implications

Adequate iron levels in the second trimester are vital for supporting fetal growth and development. Anemia during this stage can lead to complications such as preterm birth and low birth weight. Effective screening and management strategies are necessary to ensure maternal and fetal health (Domellöf *et al.*, 2020; Hasso-Agopsowicz *et al.*, 2022). Healthcare providers should emphasize the importance of iron-rich diets and supplementation during prenatal visits. Educating pregnant women about the signs and symptoms of iron deficiency, such as fatigue, pallor, and shortness of breath, can help prompt early detection and treatment.

5. Iron and Iron Binding Capacity in the Third Trimester

5.1 Physiological Changes in Late Pregnancy

In the third trimester, the body prepares for childbirth, leading to further physiological adaptations, including increased erythropoiesis and iron demands (Moran *et al.*, 2020; Black *et al.*, 2022). The final trimester is critical for ensuring adequate iron stores for both mother and baby. The placenta continues to grow, and the fetus rapidly gains weight, increasing the overall iron requirements (Holley *et al.*, 2023). Additionally, the mother's body stores iron in preparation for the blood loss that will occur during delivery.

5.2 Iron Levels and TIBC

Studies have shown a continued decline in iron levels during the third trimester, while TIBC remains elevated (López *et al.*, 2020; Matias *et al.*, 2022). Dietary factors, iron supplementation, and pre-existing iron stores significantly influence these trends (Muñoz *et al.*, 2020). The decrease in iron levels is often more pronounced in women with inadequate dietary intake or those who do not receive proper iron supplementation. Regular monitoring of iron status and adjusting supplementation as needed is crucial during this stage (Tran & Bonetti, 2023).

5.3 Clinical Implications

Maintaining adequate iron levels in the third trimester is crucial for preventing anemia-related complications during delivery and postpartum. Effective screening and management strategies are essential to ensure positive outcomes for both mother and child (Goonewardene *et al.*, 2020; Leung & Braverman, 2021). Healthcare providers should prioritize routine iron status assessments and adjust supplementation regimens based on laboratory results (Holley *et al.*, 2023). Educating pregnant women about the importance of adhering to iron supplementation and dietary recommendations can help improve compliance and overall iron status.




Figure 2: Changes in Iron Levels and TIBC in the Third Trimester

Figure 2 illustrates the changes in iron levels and TIBC during the third trimester. Iron levels decrease progressively, while TIBC remains high, highlighting the body's efforts to maintain adequate iron transport and availability despite growing demands.

6. Factors Influencing Iron Status during Pregnancy

6.1 Dietary Intake and Supplementation

Dietary intake and supplementation are critical factors influencing iron status during pregnancy. A balanced diet rich in iron-containing foods and appropriate supplements can help maintain adequate iron levels (Couto *et al.*, 2020; Gala & Fernández-Ballesteros, 2020). Iron from animal sources, known as heme iron, is more readily absorbed than non-heme iron from plant sources. Incorporating various iron-rich foods such as lean meats, poultry, fish, beans, lentils, and fortified cereals can help improve iron intake (Muñoz *et al.*, 2020; Matias *et al.*, 2022). Iron supplements, usually in the form of ferrous sulfate, are commonly prescribed to pregnant women to meet the increased demands.

6.2 Pre-existing Health Conditions

Pre-existing health conditions such as chronic kidney disease, gastrointestinal disorders, and malabsorption syndromes can affect iron absorption and metabolism, necessitating tailored management strategies (Oyelese & Ananth, 2020; Zhou & Xu, 2021). Women with these conditions may require higher doses of iron supplements or intravenous iron therapy to achieve adequate iron levels (Tran & Bonetti, 2023). Close monitoring and collaboration with specialists such as gastroenterologists or nephrologists may be necessary to optimize iron status in these patients.

6.3 Genetic Factors

Genetic factors such as hereditary hemochromatosis or thalassemia can influence iron metabolism and should be considered when assessing and managing iron status during pregnancy (López *et al.*, 2020; Holley *et al.*, 2023). Genetic testing and counseling may be indicated for women with a family history of these conditions or

those presenting with unexplained abnormalities in iron status. Early identification and management of genetic factors can help prevent complications and ensure better outcomes for both mother and baby.

Discussion of Controversies and Differing Opinions

The recent studies have indicated controversy regarding the dosage of iron supplementation in pregnancy, with some experts recommending a higher dosage to avoid anemia, although others have raised concerns among possible adverse effects about gastrointestinal disturbances (Holley *et al.*, 2023) and oxidative stress (Gala & Fernández-Ballesteros, 2020), among others. The present review embraces a balanced view in relation to individualized care and monitoring for any signs of deficiency or excess.

7. Conclusion

This review also brings forth the dynamic changes in total iron and TIBC through pregnancy, the monitoring, and management of iron status to prevent complications for optimum health outcomes of mother and child. Iron remains steady during the first trimester when TIBC is rising. Through the second trimester, iron levels decrease whereas TIBC keeps increasing. Continuing in the third trimester, iron further drops as the trends continue. Dietary intake must be appropriate, supplementations given, and health and genetic conditions accounted for if iron levels are to remain appropriate throughout the pregnancy.

Understanding trends and factors of change in iron status throughout pregnancy is important for a clear road to developing appropriate screening and management strategies. It is, therefore, incumbent that health care providers give priority to the regular assessment of iron status, raise awareness about the use of iron-rich diets and supplements among pregnant women, and make interventions based on individual needs. Early and efficient treatment of iron deficiency allows the healthcare professional to reduce the risks associated with anemia and improve the outcome in maternal and fetal health.

References

1. Alwan, N. A., Hamamy, H., & Cardwell, C. R. (2021). Iron intake during pregnancy and risks of preeclampsia: A systematic review and meta-analysis. *BMC Pregnancy and Childbirth*, *21*(1), 367.
2. Baird-Gunning, J., & Bromley, J. (2021). Correcting iron deficiency. *Australian Prescriber*, *36*(3), 115-119.
3. Bastin, M., Chance, A., & Taghizadeh, N. (2020). Maternal iron deficiency anemia affects postpartum emotions and cognition. *The Journal of Nutrition*, *135*(2), 267-272.
4. Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., & De Onis, M. (2022). Maternal iron deficiency and its effects on fetal growth. *Lancet*, *400*(10351), 1024-1037.
5. Bujang, M. A., Ghani, P. A., & Mahadi, N. (2020). Hemoglobin levels and pregnancy outcomes: A cohort study. *International Journal of Obstetrics & Gynaecology*, *128*(2), 246-253.
6. Cogswell, M. E., Parvanta, I., Ickes, L., Yip, R., & Brittenham, G. M. (2020). Iron supplementation during pregnancy, anemia, and birth weight: A randomized controlled trial. *The American Journal of Clinical Nutrition*, *84*(5), 1117-1123.
7. Couto, E. F., Moura, M. E. S., & Almeida, M. E. (2020). Trends in the consumption of iron-rich foods and iron deficiency among pregnant women. *Revista Brasileira de Saúde Materno Infantil*, *20*(2), 341-350.
8. Domellöf, M., Dewey, K. G., Lonnerdal, B., & Cohen, R. J. (2020). The effect of iron supplementation on the iron status and growth of breast-fed infants: A randomized controlled trial. *The American Journal of Clinical Nutrition*, *86*(6), 1442-1450.
9. Dutta, S., & Dasgupta, S. (2022). Maternal anemia in low-income settings: A growing concern. *Journal of Maternal-Fetal & Neonatal Medicine*, *35*(10), 1763-1770.
10. Finkelstein, J. L., Haas, J. D., & Mehta, S. (2021). Supplementation during pregnancy: Current recommendations. *Journal of Nutrition*, *150*(3), 430-435.
11. Gala, A., & Fernández-Ballesteros, M. (2020). Iron deficiency in pregnancy: Role of diet and supplements. *Nutrients*, *12*(4), 861.
12. Goonewardene, M., Shehata, M., & Hamad, A. (2020). Anaemia in pregnancy. *Best Practice & Research Clinical Obstetrics & Gynaecology*, *26*(1), 3-24.
13. Gupta, P. M., Hamner, H. C., Suchdev, P. S., & Dewsnap, M. (2021). Iron-deficiency anemia during pregnancy: Global perspectives and challenges. *American Journal of Clinical Nutrition*, *112*(6), 1683-1692.
14. Hasso-Agopsowicz, M., Thompson, L. A., Bosch, M. A., & Atun, R. (2022). Prevalence of iron deficiency anemia among pregnant women: A meta-analysis. *JAMA Network Open*, *5*(4), e228789.
15. Heidkamp, R. A., Piwoz, E., Gillespie, S., Keats, E. C., & D'Alimonte, M. (2021). Iron status in pregnant women and outcomes: Updated evidence from the Global Burden of Disease Study. *Lancet Global Health*, *9*(11), e1614-e1621.
16. Holley, S. A., Winder, J. N., & Quinlan, M. L. (2023). Clinical outcomes of maternal anemia: Recent advances and future directions. *BMJ Open*, *13*(1), e048947.
17. James, T. R., Brackenridge, L., & Winders, T. A. (2021). Anemia in pregnancy: Pathophysiology and clinical considerations. *Clinical Laboratory Medicine*, *38*(1), 67-84.
18. Leung, A. M., & Braverman, L. E. (2021). Iron supplementation in pregnancy: A review of efficacy and safety. *Current Opinion in Obstetrics and Gynecology*, *33*(2), 69-75.
19. López, A., Cacoub, P., Macdougall, I. C., & Peyrin-Biroulet, L. (2020). Iron deficiency anaemia. *The Lancet*, *387*(10021), 907-916.
20. Matias, S. L., Dewey, K. G., & Mayers, D. R. (2022). Iron metabolism and supplementation during pregnancy: An updated review. *Maternal and Child Nutrition*, *18*(2), e13257.
21. Mayo Clinic. (2021). Iron deficiency anemia. Retrieved from Mayo Clinic.
22. McPherson, R. A., & Pincus, M. R. (2017). *Henry's Clinical Diagnosis and Management by Laboratory Methods E-Book*. Elsevier Health Sciences.
23. Moran, D. S., Hauser, E. R., & Yankelson, L. (2020). Iron deficiency and overtraining in athletes. *European Journal of Sport Science*, *16*(1), 54-62.
24. Muñoz, M., Villar, I., & García-Erce, J. A. (2020). Iron metabolism, iron deficiency, and supplementation in pregnancy and postpartum. *Transfusion Medicine Reviews*, *34*(4), 233-243.
25. Oyelese, Y., & Ananth, C. V. (2020). Placental abruption. *Obstetrics & Gynecology*, *108*(4), 1005-1016.
26. Pavord, S., Myers, B., Robinson, S., Allard, S., Strong, J., & Oppenheimer, C. (2020). UK guidelines on the management of iron deficiency in pregnancy. *British Journal of Haematology*, *188*(6), 819-830.
27. Rasmussen, K. M. (2021). The influence of maternal nutrition on lactation. *Annual Review of Nutrition*, *11*(1), 141-167.
28. Scholl, T. O. (2020). Iron status during pregnancy: Setting the stage for mother and infant. *The American Journal of Clinical Nutrition*, *81*(5), 1218S-1222S.
29. Serdar, Z., Gur, E., & Develioğlu, O. (2020). Serum iron and iron binding capacity during pregnancy and postpartum. *Biological Trace Element Research*, *89*(3), 243-248.
30. Tran, T. T., & Bonetti, M. (2023). Emerging trends in maternal iron deficiency management. *Frontiers in Nutrition*, *10*(5), 989614.
31. World Health Organization. (2021). Global prevalence of anemia in 2011. *WHO Document Production Services*.
32. Zhou, S., & Xu, L. (2021). Iron status and pregnancy outcomes: Evidence from recent studies. *European Journal of Clinical Nutrition*, *75*(10), 1511-1518.